

1,3-BIS(2-HYDROXYIMINOMETHYLPYRIDINIUM) PROPANE AS THE POTENTIAL REACTIVATOR OF THE ACETYLCHOLINESTERASE INHIBITED BY NERVE AGENTS

Kamil Kuča¹, Vlastimil Dohnal²

Purkyně Medical Military Academy in Hradec Králové, Czech Republic: Department of Toxicology¹; Masaryk University in Brno, Faculty of Science, Czech Republic: Department of Analytic Chemistry²

Summary: The oxime K005 [1,3-bis(2-hydroxyiminomethylpyridinium) propane dibromide] for the reactivation of the enzyme acetylcholinesterase (AChE) inhibited by cyclosarin and VX was tested. Its reactivation potency was not better in comparison with the currently used AChE reactivator - pralidoxime. On the other hand, the oxime K005 has its maximum reactivation ability at the concentration 10^{-4} M, which could be achieved for human use. The maximum concentration of pralidoxime was reached at the concentrations 10^{-1} M (for cyclosarin) and 10^{-3} M (for VX) but these concentrations are not available for the use in vivo.

Key words: Acetylcholinesterase; Reactivation; Inhibition; Oximes; Reactivator; Pralidoxime; Cyclosarin; VX; In vitro; K005; Nerve agents; Organophosphates

Introduction

Acetylcholinesterase (AChE; EC 3.1.1.7) reactivators and parasympatolytics are currently used as the antidotes for the treatment of the organophosphate compounds (OPCs) intoxications (10). OPCs are largely used in agriculture (insecticides), in industry and technology (softening agents) and in military technology (chemical warfare agents) (4). OPCs affect AChE as irreversible inhibitors. They phosphorylate or phosphonylate the serine hydroxyl group at the active site of the enzyme (6). Due to the inhibition, AChE cannot be physiologically functional (10).

Pralidoxime [1-methyl-2-hydroxyiminomethylpyridinium chloride], obidoxime [1,3-bis(4-hydroxyiminomethylpyridinium)-2-oxa-propane dichloride] and H-oxime HI-6 [1-(2-hydroxyiminomethylpyridinium)-3-(4-carbamoylpyridinium)-2-oxa-propane dichloride] are currently considered to be the main representatives of the AChE reactivators (5). These compounds have in their molecules incorporated characteristic features: the presence of the quaternary nitrogen (in the case of obidoxime and HI-6 even two quaternary nitrogen atoms), functional aldoxime group and in the case of bisquaternary pyridinium aldoximes, the connecting chain of the specific length and shape (9).

None from the above mentioned AChE reactivators are able to satisfactorily reactivate AChE inhibited by all kinds

of OPCs, especially by nerve agents such as tabun, sarin or cyclosarin (5).

The reactivation insufficiency is caused by the differences in the structure of the AChE inhibitors (3). According to the insufficient reactivation potency of all currently used reactivators, scientists in many laboratories throughout the world synthesize and test new AChE reactivators, with the aim to treat intoxications by all known OPCs (11).

In our Department we have synthesized 1,3-bis(2-hydroxyiminomethylpyridinium) propane dibromide (called K005) as the potential reactivator of the AChE inhibited by nerve agents (7). In this work we have tested the potency to reactivate AChE inhibited by highly toxic nerve agents such as cyclosarin and VX.

Material and Methods

Chemical

Compound K005 [1,3-bis(2-hydroxyiminomethylpyridinium) propane dibromide] was prepared earlier at our department (7) by the modification of the synthesis described by Berry et al. (1). Pralidoxime [1-methyl-2-hydroxyiminomethylpyridinium chloride] was used for comparison as the currently used AChE reactivator. Purities of the tested AChE reactivators were detected using HPLC. Nerve agents (cyclosarin and VX) were obtained from the Military Facility Brno (97% purity). All other chemicals used were of reagent grade (Aldrich).

Enzyme

Rat brain acetylcholinesterase was chosen as the source of the enzyme. Its preparation was as follows. Lightly ether-narcotized animals were killed by bleeding from a carotid artery and the brains were removed, washed with saline and homogenized in an Ultra-Turrax homogenizer in a distilled water to make a 10 % homogenate.

Tab. 1: Kinetic parameters for the reactivation of cyclosarin inhibited AChE.

| Oxime | K_R [μM] | k_R [min^{-1}] | k_r [$\text{min}^{-1}\text{M}^{-1}$] |
|-------|-------------------------|-----------------------------|--|
| 2-PAM | 12000 | 0.040 | 3 |
| K005 | 5 | 0.010 | 2196 |

Tab. 2: Kinetics parameters for the reactivation of VX inhibited AChE.

| Oxime | K_R [μM] | k_R [min^{-1}] | k_r [$\text{min}^{-1}\text{M}^{-1}$] |
|-------|-------------------------|-----------------------------|--|
| 2-PAM | 127 | 0.047 | 370 |
| K005 | 1350 | 0.132 | 98 |




Fig. 1: Concentration-reativation relationship for the reactivation of cyclosarin inhibited AChE.




Fig. 2: Concentration-reativation relationship for the reactivation of VX inhibited AChE.

In vitro measurement

Reactivation effectivities of the oximes were tested *in vitro* on the model of AChE inhibited by cyclosarin using standard reactivation test with electrometric instrumentation (8). The AChE homogenate (0.5 ml) was mixed with 0.5 ml of 0.01 μM cyclosarin in dry isopropanol and incubated for 30 min (25 °C). Then 2.5 ml of 3M NaCl was added and supplied by distilled water to a volume of 23 ml. After that, 2 ml of 0.02 M acetylcholine bromide was added and enzyme activity was assayed titrimetrically at pH 8.0 and 25 °C on the Autotitrator RTS 822 (Radiometer, Denmark).

The activities of intact (a_o) and GF-inhibited (a_i) AChE were determined. When GF-inhibited AChE was incubated 10 min with solution of reactivator, the activity of reactivated AChE (a_r) was obtained. The activity values a_o , a_i and a_r were calculated from the slopes of the initial part of titration curves. Each value represents arithmetic mean from two independent measurements.

Results

Affinity to the intact enzyme

Constant K_{DIS} characterizes the affinity of the AChE reactivators to the intact enzyme. The lower is its value, the higher is its affinity to the intact enzyme. In our case, compound K005 has value of the dissociation constant four times lower (53 μM) compared to pralidoxime (210 μM). Thus, its affinity to the intact enzyme is higher.

Reactivation of cyclosarin inhibited AChE

The results obtained for the cyclosarin inhibited AChE are summarized in Table 1. Values of the constant k_R (characterizing the decomposition of the inhibited enzyme-reactivator complex) and k_r (characterizing velocity of overall reaction) favour compound K005. On the other hand, K_R (characterizing the affinity of the enzyme to the inhibited AChE) is advantageous for the pralidoxime. The relationship between the concentration of the oxime (concentration range from 10^{-8} to 10^{-1} M) and reactivation potency of the oximes is shown in Figure 1. The concentration with the maximum reactivation potency for the oxime K005 is at the concentration 10^{-4} M (10 %). On the other hand, pralidoxime is able to reactivate cyclosarin-inhibited AChE *in vitro* in very high concentrations, and therefore, it would be toxic for the use *in vivo*.

Reactivation of VX inhibited AChE

Kinetics parameters characterizing the ability of the oxime K005 and pralidoxime to reactivate VX-inhibited AChE are summarized in Table 2. Both reactivators were able to reactivate VX-inhibited AChE. Value of the constant K_R characterizing affinity to the inhibited enzyme favours pralidoxime. Its affinity to the inhibited enzyme is almost 11 times higher compared to the oxime K005. Constant k_R characterizing the decomposition of the inhibited enzyme-reactivator complex is about three times higher for the oxime

me K005. On the other hand, constant k_r , characterizing the velocity of the overall reaction favours pralidoxime and its value is about 3.5 times higher compared to oxime K005.

The concentration-reactivation relationship is demonstrated in the Figure 2. 10^{-3} M concentration of the pralidoxime is necessary to reach 34 % reactivation of VX inhibited AChE. In the case of the oxime K005, maximum reactivation potency (6.35 %) is reached at the concentration 10^{-4} M.

Discussion

Oxime K005 does not reach better reactivation potency than pralidoxime in the case of cyclosarin and VX-inhibited AChE. On the other hand, concentration with maximum reactivation potency was reached at the concentration lower than that in the case of the pralidoxime. These concentrations could be safe for human use (8).

Reactivation of inhibited AChE depends on the inhibitors used and on the chemical structure of the reactivators (3,5,8). Our results confirm these observations. There are differences in the course of the reactivation curves of the oxime K005 or pralidoxime for inhibition by cyclosarin or VX.

Shape of the molecules of the AChE reactivators affects reactivation process, too (2,9). According to our results, there were significant differences in the reactivation potency between oxime K005 and pralidoxime. Pralidoxime has in its molecule one quaternary nitrogen only while oxime K005 has two quaternary nitrogens. The presence of the quaternary nitrogen in the molecule is an important factor for reactivation process (12).

In conclusion, we have tested oxime K005 [1,3-bis(2-hydroxyiminomethylpyridinium) propane dibromide] for the reactivation of the enzyme AChE inhibited by cyclosarin and VX. Its reactivation potency was not better in comparison with currently used AChE reactivator – pralidoxime. On the other hand, oxime K005 has maximum reactivation ability at the concentration 10^{-4} M, which could be safe for human use. Maximum concentration of pralidoxime was reached at the concentrations 10^{-1} M (for

cyclosarin) and 10^{-3} M (for VX). Nevertheless, these concentrations are not available for the use in vivo.

Acknowledgement

The authors are grateful to Mrs. I. Ježková and Mrs. M. Hrabínová for their skilful technical assistance. The study was supported by the grant of Ministry of Defense No. ONVLAJEP20031.

References

1. Berry WK, Davies DR, Green AL. Oximes of α,ω -diquaternary alkane salts as antidotes to organophosphate anticholinesterases. *Brit J Pharmacol* 1959;14: 186–91.
2. Bielavský J, Kassa J, Elsnerová E, Dejmeck L. Cholinesterase reactivators derived from pyridine-2-carbaldoxime. *Collect Czech Chem Commun* 1997;63:199–204.
3. Cabal J, Kassa J, Patočka J. Inhibition of plasma cholinesterase by O-alkyl-fluorophosphonates. *Collect Czech Chem Commun* 1996;62:521–6.
4. Kassa J. Review of oximes in the antidotal treatment of poisoning by organophosphorus nerve agents. *J Toxicol Clin Toxicol* 2002;40:803–16.
5. Kuča K, Bielavský J, Cabal J, Bielavská M. Synthesis of a potential reactivator of acetylcholinesterase – 1-(4-hydroxyiminomethylpyridinium)-3-(carbamoylpyridinium)-propane dibromide. *Tetrahedron Lett* 2003;44:3123–5.
6. Kuča K, Bielavský J, Cabal J, Kassa J. Synthesis of a New Reactivator of tabun-inhibited acetylcholinesterase. *Bioorg Med Chem Lett* 2003;13:3545–7.
7. Kuča K, Cabal J, Patočka J, Kassa J. Synthesis of bisquaternary symmetric – χ,δ -bis(2-hydroxyiminomethylpyridinium)alkane dibromides and their reactivation of cyclosarin-inhibited acetylcholinesterase. *Lett Org Chem* 2004;1:84–6.
8. Kuča K, Kassa J. A comparison of the ability of a new bispyridinium oxime – 1-(4-hydroxyiminomethylpyridinium)-4-(4-carbamoylpyridinium)butane dibromide and currently used oximes to reactivate nerve agent-inhibited rat brain acetylcholinesterase by in vitro methods. *J Enzyme Inhib Med Chem* 2003;18:529–35.
9. Kuča K, Patočka J, Cabal J. Reactivation of organophosphate inhibited acetylcholinesterase activity by α,ω -bis-(4-hydroxyiminomethylpyridinium)alkanes in vitro. *J Appl Biomed* 2003;1:207–11.
10. Marrs TC. Organophosphate poisoning. *Pharmacol Therap* 1993;58:51–66.
11. Petrova I, Bielavský J. An overview of syntheses of cholinesterase reactivators from 1980 to 1992. *Voj Zdrav Listy* 2001;70:63–73 (in Czech).
12. Ševčíková K, Hampl F, Liška F, Cabal J, Patočka J. Substituted pyridinium derivatives as inhibitors of acetylcholinesterase. *Voj Zdrav Listy – Supplementum* 1998;67:28–31.

Submitted October 2003.

Accepted June 2004.

*Ing. Kamil Kuča,
Purkyně Medical Military Academy,
Department of Toxicology,
Třebešská 1575, 500 01 Hradec Králové,
Czech Republic.
e-mail: kucakam@pmfhk.cz*